

Do clusters of galaxies rotate?

Hrant M. Tovmassian

*Instituto Nacional de Astrofísica Óptica y Electrónica, AP 51 y 216, 72000,
Puebla, Pue, México*

September 2002

Abstract. It is shown that the number of members, N_A , and the linear radius of the Abell clusters at $z < 0.18$ increase with redshift. The increase is caused by observational selection: the number of rich and large clusters is small in the nearby small volume of space, while poor and small clusters are not detected at high distances. Besides, the numbers of clusters with different ellipticities are about the same at all distances, though in the case of cosmological evolution the number of the more elliptical clusters increases with decrease of distance. It means that clusters do not evolve dynamically. The dependences of the X-ray emission detection rate and the temperature of the X-ray emission on redshift are consequences of the dependence of N_A on redshift. The radial velocity dispersion of the less elliptical clusters is higher than that of the elongated ones. This shows that galaxies in a cluster move preferentially along the large axis of the cluster. The space density is not changing in the considered range of z . Movement of galaxies along the cluster elongation may be explained by rotation along the large axis of a cluster.

Keywords: galaxies: clusters: general – dynamics: galaxies

It is supposed that clusters grow by accretion and merging of galaxies along filamentary large-scale structures (West 1994, West et al. 1995). Plionis (2002) claims that nearby clusters at $z < 0.15$ are still evolving. It means, that nearby clusters should be richer and perhaps larger than the distant ones. I compared the number N_A of galaxies (Struble & Rood 1987) in ACO clusters (Abell et al. 1989) and their linear radii R_l (in Mpc) with redshift z (Figs. 1 and 2). For analysis I used 621 clusters with z determined by at least 2 galaxies (Struble & Rood 1999) and located at $z < 0.18$. Figs. 1 and 2 show that N_A and R_l increase with z . The dashed regression lines in both Figures are drawn after exclusion of 381 clusters of the richness class 0, the sample of which is assumed to be incomplete. The Spearman rank test showed that the correlation factors for the total sample and for the sample without richness class 0 clusters are $\rho = 0.41$ and $\rho = 0.42$ with significances $\alpha = 10^{-27}$ and $\alpha = 10^{-12}$ respectively. Corresponding values for R_l (Fig. 2) are $\rho = 0.44$ and $\rho = 0.37$, and $\alpha = 10^{-31}$ and $\alpha = 10^{-9}$ respectively. Since exclusion of the richness class 0 clusters does not alter these dependences, in the further analysis I use the whole sample of clusters. Apparently the dependences of N_A and R_l on z are due to a selection effect. At high z mostly rich and large clusters are detected. The number of them is small in the nearby small volume of space. Poor



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.

and small clusters located at high z are missed in the background. If dependences of N_A and R_l on z are caused by evolution, N_A and R_l should be correlated. However, comparison shows no correlation ($\rho = 0.119$, $\alpha = 0.003$) between N_A and R_l .




Figure 1. The dependence of N_A on z . The solid line is the regression line determined using the data on 616 clusters. The dashed line is determined for a sample in which 378 poor clusters of richness 0 are excluded.




Figure 2. The dependence of the linear R_c on z . The solid and dashed lines are determined as in Fig. 1.

Besides, in the case of cosmological evolution the number of elliptical clusters (ECs) should be higher at high z , and the number of round clusters (RCs) should be high at small z (Mellot et al. 2001). I compared the distribution of the number of ECs with $f \leq 1.18$ ¹ with that of the RCs with $f \geq 2.0$ (Fig. 3). The values of f are taken from Struble & Ftaclas (1995). The K-S test showed that the hypothesis that two distributions are of different population is rejected with probability 0.6, i.e. both samples are consistent with a single distribution function. Moreover, the RCs should accumulate in time more members and should be more rich than ECs. Meanwhile, the mean number of mem-

¹ $f = a/b$ where a is the length of the cluster and b is its width.

bers of 18 RCs is 61 ± 22 , and does not differ from that of the 27 ECs, being equal to 63 ± 32 . In the case of cosmological evolution the cluster velocity dispersion, σ_v , should increase in time since virialization will tend to increase the cluster “thermal” velocity dispersion (e.g. Plionis 2002). I considered σ_v ’s of a sample of 362 clusters with σ_v determined by at least five galaxies (Struble & Rood 1999). The Spearman rank test showed that, contrary to expectation, σ_v is not correlated with z ($\rho = 0.128$ and the significance $\alpha = 0.0146$).




Figure 3. The distribution of the numbers N of the RCs (solid line) and ECs (dashed line) by z .

If clusters grow by accretion of the surrounding matter, then the space density ρ of galaxies should increase in time, i.e. it should be higher in the nearby space. Fig. 4 shows that ρ increases by 4.5 times from $z = 0.18$ to $z = 0.02$. However, just by this amount ρ must increase due to increase of N_A and R_l with z (Figs. 1 and 2) which, in turn, is a result of observational selection.

Hence, observational data *contradict to cosmological evolution* of clusters up to $z = 0.18$.

The observed dependence of N_A on z may cause correlation between other cluster parameters. In Table 1 the average N_A of clusters with detected X-ray emission is compared with that of clusters with no X-ray detection (Ebeling et al. 1996, 1998, 2000). It is seen that in each





Figure 4. The dependence of the space density of galaxies in clusters on z .

Table I. The average numbers N_A of clusters with and without X-ray emission.

z	< 0.06	$0.06-0.10$	$0.10-0.14$	$0.14-0.18$
D-ted	54.7 ± 19.4 (57)	63.4 ± 21.7 (65)	90.8 ± 33.2 (23)	131.2 ± 64.7 (10)
Not d-ted	46.5 ± 14.2 (69)	54.1 ± 25.5 (191)	64.3 ± 27.7 (128)	84.7 ± 36.0 (70)

Figure 5. The dependence of kT on N_A .

of the four ranges of z the average N_A of clusters with X-ray emission is higher than in clusters without X-ray emission. In Fig. 5 the ICM gas temperature T of 213 clusters is compared with N_A . It shows that T increases with N_A : $\log kT \sim 0.54 \log N_A$. The correlation factor $\rho = 0.54$ and significance $\alpha = 1.7 \cdot 10^{-17}$, i.e the correlation is strong. On the other hand, it has been shown that the temperature T of the X-ray emitting gas and its mass M_{gas} are correlated: $M_{gas} \sim T^{1.5}$ (Evrard et al. 1996) or $M_{gas} \sim T^{1.79}$ (Nevalainen et al. 2000) at $z = 0$. It follows that $M_{gas} \sim N_A^{0.96}$, i.e. the ISM mass is strictly proportional to the number of galaxies. Thus, the correlations of the X-ray detection rate and the X-ray temperature T on z are simply consequences of the dependence of N_A on z , caused by observational selection.

Tovmassian (2001, 2002) showed that the radial velocity dispersion σ_v of compact groups (CGs) is minimal for elongated chain-like groups with small b/a ratio, while the less elongated groups have higher σ_v . It shows that the member galaxies in a group move preferentially along the elongation of a group. The correlation σ_v versus b/a holds in the case of clusters as well. The mean σ_v of 15 RCs is 917 ± 338 , and that of 18 ECs is 642 ± 254 . The t test shows that the difference is highly significant ($\alpha = 0.012$). The mean N_A of 12 RCs is 62 ± 17 , while that of 17 ECs is 60 ± 25 . It means that the difference in σ_v is not caused by the difference in N_A . Therefore, we conclude, that the observed difference of σ_v in RCs and ECs is due to a *projection* effect. Hence, galaxies in clusters, as in CGs, move *preferentially along the large axis*. In clusters

the large axis of which is oriented close to the line of sight and which are seen more round, a large number of member galaxies move towards us or backwards. In such clusters σ_v is high, and is close to the real value. The σ_v of ECs oriented, on average, at $\sim 45^\circ$ to the line of sight should be by $\approx \sin 45^\circ$ smaller than those of RCs, i.e. should be equal to 648 km s^{-1} . The mean σ_v of ECs differs only by $\approx 10\%$ from the estimated value.

Movement of galaxies along the large axis of a cluster may be explained by three ways. The option that clusters grow by accretion and merging of matter mainly along the large axis (West 1994, West et al. 1995) is hardly valid, since as we showed, observational data contradict to this option. Moreover, in the case of accretion only a few infalling galaxies would move along the cluster elongation, but not the majority of its members. Another option is departure of member galaxies from each other in opposite directions. The reason may probably be the influence of another clusters. However, the velocities initiated by the gravity of clusters with masses $\sim 10^{14}$ located at opposite directions at distances of $\approx 30 \text{ Mpc}$ are $\approx 30 \text{ km s}^{-1}$, and may not account for the observed difference of σ_v . No physical mechanism is yet known for explanation of expansion of clusters according to Ambartsumian (1961) or Arp (2001, and references therein) ideas. Hence, the more realistic option is *rotation of galaxies* around their gravitational center, as in the case of CGs (Tovmassian 2001, 2002).

References

- Abell, G.O., Corwin, H.C., Olowin, R., 1989, ApJS, 70, 1
 Ambartsumian V.A. 1961, AJ, 66, 536
 Arp H. 2001, ApJ, 549, 780
 Ebeling, H., Voges, W., Böhringer, H., et al., 1996, MNRAS, 281, 799
 Ebeling, H., Edge, A. C., Böhringer, H., 1998, MNRAS, 301, 881
 Ebeling, H., Edge, A.C., Allen, S.W., et al., 2000, MNRAS, 318, 333
 Melott, A.L., Chambers, S.W., Miller, C.J. 2001, ApJ, L75
 Nevalainen, J., Markevitch, M., Forman, W. 2000, ApJ, 532, 694
 Plionis, M. 2002, ApJ, 572, 67L
 Struble, M. F., Ftaclas, C., 1995, AJ, 108, 1
 Struble, M. F., Rood, H. J., 1987, ApJS, 63, 555
 Struble, M. F., Rood, H. J., 1999, ApJS, 125, 35
 Tovmassian, H. M., 2001, PASP, 113, 543
 Tovmassian H. M., 2002, AN, 323, 488
 West, M. J. 1994, MNRAS, 268, 79
 West, M.J., Jones, C., Forman, W. 1995, ApJ, 451, L5

